

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C12Q 1/68, C12P 19/34, C12N 15/11, 15/85, C07H 21/04</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/61810 (43) International Publication Date: 19 October 2000 (19.10.00)</p>
<p>(21) International Application Number: PCT/US00/09293 (22) International Filing Date: 7 April 2000 (07.04.00) (30) Priority Data: 60/128,377 8 April 1999 (08.04.99) US (71) Applicant (for all designated States except US): OASIS BIO-SCIENCES, INC. [US/US]; 3550 General Atomics Court, San Diego, CA 92121-1122 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BROWN, Bob, D. [US/US]; 445 North Willow Spring Drive, Encinitas, CA 92024 (US). RILEY, Timothy, A. [US/US]; 12894 War Horse Street, San Diego, CA 92129 (US). (74) Agent: ISRAELSEN, Ned, A.; 16th floor, 620 Newport Center Drive, Newport Beach, CA 92660 (US).</p>	<p>(81) Designated States: AE, AG, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, DZ, EE, ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KR (Utility model), KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	
<p>(54) Title: ANTISENSE OLIGONUCLEOTIDES COMPRISING UNIVERSAL AND/OR DEGENERATE BASES</p> <p>bcl-2A: 2020 tggatccaggataacggaggctgggatgcctttgtggaact 2060 * * * bcl-xL: 675 tggatccaggagaacggcggtgggatacttttgtggaact 715 Examples: 3' ACCTAGGTCCTBTTGCCBCC 3' GGTCCTBTTGCCBCCGACCC 3' CCTBTTGCCBCCGACCCTAPGKAAACACC 3' GCCBCCGACCCTAPGKAAAC 3' BCCGACCCTAPGK 3' ACCCTAPGKAAACACCTTGA</p>		
<p>(57) Abstract</p> <p>Antisense oligonucleotides containing one or more degenerate and/or universal bases, and one or more modified backbone linkages, and use of these oligonucleotides for cleaving target RNA molecules.</p>		

062

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ANTISENSE OLIGONUCLEOTIDES COMPRISING UNIVERSAL AND/OR DEGENERATE BASESField of the Invention

The present invention relates to antisense oligonucleotide compositions comprising one or more universal
5 and/or degenerate bases, and to methods for using these oligonucleotides to target RNA molecules

Description of the Related Art

Antisense technology is based on the finding that gene expression can be modulated using an oligonucleotide
which binds to the target RNA. By exploiting the Watson-Crick base pairing and the ability to recruit certain
nucleases, particularly RNase H, to specifically cleave the target RNA in the DNA/RNA hybrid, one can design
10 antisense molecules which are highly specific for the target nucleic acid molecule. However, there are families of
genes in which this high degree of specificity may be detrimental. For example, it may be desirable to target two or
more of these genes if there is a synergistic effect if the genes are inactivated together.

Typical antisense compounds are modified nucleic acids that bind to their target RNA via Watson-Crick base
pairing. Different constructions can recruit a variety of RNases to mediate the cleavage of the target RNA. The most
15 common RNase is RNase H which recognizes a DNA/RNA duplex, followed by cleavage of the target RNA. The
oligonucleotide most commonly used for this purpose contains unmodified (naturally-occurring) bases (A, T, G, C) and a
modified backbone called a phosphorothioate which renders the oligonucleotide resistant to nucleases. Other backbone
modifications such as 2'-O-alkyl render the oligonucleotide unable to mediate RNase H cleavage of the target RNA.
There are many reports of the combination of non-RNase H substrate portions and RNase H substrate portions within a
20 single antisense oligonucleotide. These non-RNase H substrate portions provide both binding and specificity for the
antisense oligonucleotide. Examples of these backbones include methylphosphonates, morpholinos, MMI, peptide
nucleic acids (PNA) and 3'-amidates. Sugar modifications that increase antisense oligonucleotide binding and nuclease
stability include 2'-O-alkyl, 2'-O-allyl, 2'-O-methoxyethyl, 2'-O-alkylaminoalkyl, 2'-fluoro (2'-F) and 2'-amino.

Universal or degenerate bases are heterocyclic moieties which have the ability to hydrogen bond to more than
25 one base in a DNA duplex without destroying the ability of the whole molecule to bind to the target. The use of
oligonucleotides having unmodified backbones and containing degenerate or universal bases is known in the PCR primer
literature (Bergstrom et al., *J. Am. Chem. Soc.* 117:1201-1209, 1995; Nichols et al., *Nature* 369:492-493, 1994;
Loakes, *Nucl. Acids Res.* 22:4039-4043, 1994; Brown, *Nucl. Acids Res.* 20:5149-5152, 1992). However, to date
these universal and degenerate bases have not been used in antisense technology, and have not been incorporated into
30 oligonucleotides which comprises modified backbone linkages. The present invention addresses these antisense
compositions and methods.

Summary of the Invention

One embodiment of the present invention is an antisense oligonucleotide having at least one non-naturally
occurring backbone linkage and having between 6 and about 50 bases, wherein at least one of the bases are universal
35 and/or degenerate bases. Preferably, no more than about 50% of the bases are universal and/or degenerate bases.

Another embodiment of the present invention is an antisense oligonucleotide comprising a first non-RNase H recruiting region having between 3 and about 15 bases, an RNase H recruiting region having between 3 and about 15 bases, and a second non-RNase H recruiting region, wherein at least one of the bases are universal and/or degenerate bases. Preferably, no more than about 50% of the bases are universal and/or degenerate bases.

5 The present invention also provides an antisense oligonucleotide comprising a non-RNase H recruiting section and an RNase H recruiting section, wherein at least one but of the bases are universal and/or degenerate bases. Preferably, no more than about 50% of the bases are universal and/or degenerate bases.

Another embodiment of the present invention is an oligonucleotide comprising an RNase L-recruiting region comprising a 2'-5' adenosine oligomer, wherein at least one of the bases in the RNA targeting region of the
10 oligonucleotide are universal and/or degenerate bases. Preferably, not more than about 50% of the bases in the RNA targeting region are universal and/or degenerate bases.

The present invention also provides an oligonucleotide designed to recruit RNase P, wherein at least one of the bases in the RNA targeting region of the oligonucleotide are universal and/or degenerate bases. Preferably, no more than about 50% of the bases in the RNA targeting region are universal and/or degenerate bases.

15 Another embodiment of the present invention is a ribozyme having at least one universal and/or degenerate base in its RNA targeting region. Preferably, no more than about 50% of the bases in the RNA targeting region are degenerate and/or universal bases.

The present invention also provides a method for cleaving a target RNA molecule, comprising the step of contacting the RNA molecule with any of the oligonucleotides described above in the presence of an RNase activity
20 capable of cleaving the target. Preferably, the RNase is RNase H, RNase L or RNase P.

The present invention also provides a method for cleaving a target RNA molecule, comprising the step of contacting the RNA molecule with the ribozyme described above.

The present invention also provides a method for cleaving a target RNA molecule, comprising the step of contacting said RNA molecule with the ribozyme described above.

25 Another embodiment of the present invention is a method for cleaving a target RNA molecule, comprising the step of contacting the RNA molecule with an oligonucleotide having between 6 and about 50 bases, wherein the oligonucleotide comprises at least one universal and/or degenerate base.

The present invention also provides a method for reducing the deleterious effects of an antisense oligonucleotide comprising one or more sequence motifs, comprising replacing one or more bases within said one or
30 more sequence motifs with one or more universal and/or degenerate bases. Preferably, the sequence motif is a CG dinucleotide. In another aspect of this preferred embodiment, the sequence motif is a poly-G sequence.

Brief Description of the Drawings

Figure 1 shows a sequence alignment of a region of high homology between the human bcl-2A and human bcl-xL genes. Antisense oligonucleotides complementary to the aligned sequence region, and which include one or more
35 universal and/or degenerate bases, are shown below the sequence alignment. Base mismatches are indicated by

asterisks. B indicates a universal base. P and K are degenerate bases which pair with any pyrimidine and any purine, respectively.

Figure 2 shows a sequence alignment of three homology regions of three human protein kinase C (PKC) family members. Antisense oligonucleotides complementary to the aligned sequence region, and which include one or more universal and/or degenerate bases, are shown below the sequence alignment. These antisense oligonucleotides simultaneously target two or more PKC family members.

Figure 3 shows a sequence alignment of homology regions between two alleles of the bcl-2 gene, bcl-2B and bcl-2C. Representative antisense oligonucleotides including one or more universal and/or degenerate bases are shown below the sequence alignments.

Detailed Description of the Preferred Embodiments

The present invention provides antisense oligonucleotides including one or more universal and/or degenerate bases and methods for targeting RNA which includes a region complementary or nearly complementary to the antisense oligonucleotides. Conventional antisense oligonucleotide containing only naturally occurring nucleotide bases (A, T, G, C, and U) are efficient only when they are completely complementary to their target sequence. In other words, the oligonucleotide cannot bind with sufficient affinity to mismatched oligonucleotides. This compromises the ability of conventional oligonucleotides to bind to single nucleotide polymorphisms (SNPs), and does not permit targeting of two or more homologous genes containing one or more mismatches with a single antisense oligonucleotide. The present invention solves this problem by incorporating one or more universal and/or degenerate bases (defined below) into antisense oligonucleotides. Because these universal and/or degenerate bases can tolerate nucleotide mismatches and bind with sufficient affinity to allow recruitment of nucleases, they solve this mismatch problem.

The incorporation of at least one universal and/or degenerate base into an antisense oligonucleotide can be used to reduce or eliminate the deleterious effects caused by a series or group of natural bases. Various short base sequences in oligonucleotides cause significant sequence-dependent biological effects which are not antisense-specific. For example, almost all nucleotides containing an unmethylated "CG" dinucleotide cause a variety of immune-activation effects when injected into animals, or when incubated with isolated bone marrow cells. The most common immune activation effects are enhanced B-cell proliferation and cytokine production, including inflammatory cytokines such as interleukin-2. This immune activation phenomenon is believed to be responsible for some deleterious side effects of many therapeutic antisense oligonucleotide candidates. The present invention addresses this problem by the substitution of a degenerate or universal base for C or G in these "CG" repeats. This is believed to eliminate undesirable immune activation effects, while maintaining efficient, specific antisense activity.

In addition, "GGGG" and other poly-G motifs have been shown repeatedly to produce non-antisense effects such as growth inhibition in cell cultures and high systemic toxicity in animals. Substitution of universal and/or degenerated bases within tetra-G or other poly-G motifs can "break-up" these sequences and result in an antisense oligonucleotide having significant research and therapeutic utility in both animals and cell culture.

The term "antisense" as used herein refers to a molecule designed to interfere with gene expression and capable of recognizing or binding to a specific desired target polynucleotide sequence. Antisense molecules typically (but not necessarily) comprise an oligonucleotide or oligonucleotide analog capable of binding specifically to a target sequence present on an RNA molecule. Such binding interferes with translation by a variety of means, including preventing the action of polymerases, RNA processing and recruiting and/or activating nucleases such as RNase H, RNase L and RNase P.

The term "ribozyme" as used herein refers to an oligonucleotide or oligonucleotide analog capable of catalytically cleaving a polynucleotide.

The term "oligonucleotide" refers to a molecule consisting of DNA, RNA or DNA/RNA hybrids.

The term "oligonucleotide analog" refers to a molecule comprising an oligonucleotide-like structure, for example having a backbone and a series of bases, wherein the backbone and/or one or more of the bases can be other than the structures found in naturally-occurring DNA and RNA. "Non-natural" oligonucleotide analogs include at least one base or backbone structure that is not found in natural DNA or RNA. Exemplary oligonucleotide analogs include, but are not limited to, DNA, RNA, phosphorothioate oligonucleotides, peptide nucleic acids (PNA), methoxyethyl phosphorothioates, oligonucleotide containing deoxyinosine or deoxy 5-nitroindole, and the like.

The term "backbone" as used herein refers to a generally linear molecule capable of supporting a plurality of bases attached at defined intervals. Preferably, the backbone will support the bases in a geometry conducive to hybridization between the supported bases of a target polynucleotide.

The term "non-naturally occurring base" refers to a base other than A, C, G, T and U, and includes degenerate and universal bases as well as moieties capable of binding specifically to a natural base or to a non-naturally occurring base. Non-naturally occurring bases include, but are not limited to, propynylcytosine, propynyluridine, diaminopurine, 5-methylcytosine, 7-deazaadenosine and 7-deazaguanine.

The term "universal base" refers to a moiety that may be substituted for any base. The universal base need not contribute to hybridization, but should not significantly detract from hybridization. Exemplary universal bases include, but are not limited to, inosine, 5-nitroindole and 4-nitrobenzimidazole.

The term "degenerate base" refers to a moiety that is capable of base-pairing with either any purine, or any pyrimidine, but not both purines and pyrimidines. Exemplary degenerate bases include, but are not limited to, 6H, 8H-3,4-dihydropyrimido[4,5-c][1,2]oxazin-7-one ("P", a pyrimidine mimic) and 2-amino-6-methoxyaminopurine ("K", a purine mimic).

The term "target polynucleotide" refers to DNA, for example as found in a living cell, with which the antisense molecule is intended to bind or react.

The term "activity" refers to the ability of an antisense molecule of the invention, when hybridized to a target polynucleotide, to interfere with the transcription and/or translation of the target polynucleotide. Preferably, the interference arises because the antisense molecule, when hybridized, recruits a nuclease, and/or serves as a nuclease substrate. The term "interference" includes inhibition to any detectable degree.

The term "RNase H recruiting" refers to an oligonucleotide having at least one phosphorothioate and/or phosphodiester backbone. This type of backbone is recognized by RNase H once a RNA/DNA hybrid is formed and allows RNase H to cleave the target RNA.

The term "non-RNase H-recruiting" refers to an oligonucleotide having linkages other than deoxyphosphodiester or deoxyphosphorothioate linkages, including, but not limited to, 2'-O-alkyl, PNA, methylphosphonate, 3'-amidate, 2'-F, morpholino, 2'-O-alkylaminoalkyl and 2'-alkoxyalkyl. This type of oligonucleotide is not recognized by RNase H after formation of a DNA/RNA hybrid.

The term "RNase L recruiting" refers to an oligonucleotide comprising four consecutive adenosine bases in 2', 5'-linkage which form an oligomer. This oligomer is recognized by RNase L once a DNA/RNA hybrid is formed (See U. S. Patent No. 5,583,032).

The term "RNase P recruiting" refers to an oligonucleotide capable of forming a stem-loop structure which is recognized by RNase P, an enzyme normally involved in generation of mature tRNA by cleaving a portion of tRNA precursor molecules. This stem-loop structure resembles the native tRNA substrate and is described by Ma et al. (*Antisense Nucl. Acid Drug Dev.* 8:415-426, 1998) and in U.S. Patent No. 5,877,162.

The antisense oligonucleotides and oligonucleotide analogs of the invention are preferably between 6 and about 50 bases long, more preferably between about 10 and 30 bases long, and most preferably between about 15 and 25 bases long. Oligonucleotides having 18 base pairs are particularly preferred.

The antisense oligonucleotides and oligonucleotide analogs of the invention typically contain at least one universal or degenerate base, and at least one modified backbone linkage. In general, these oligonucleotides do not contain more than about 50% universal and/or degenerate bases.

The oligonucleotides and oligonucleotide analogs of the present invention can be synthesized using standard oligonucleotide synthesis methods (see Example 1).

The oligonucleotides used in the binding domains can employ any any backbone and any sequence capable of resulting in a molecule that hybridizes to natural DNA and/or RNA. Examples of suitable backbones include, but are not limited to, phosphodiester and deoxyphosphodiester, phosphorothioate and deoxyphosphorothioate, 2'-O-substituted phosphodiester and deoxy analogs, 2'-O-substituted phosphorothioate and deoxy analogs, morpholino, PNA (U. S. Patent No. 5,539,082), 2'-O-alkyl methylphosphonates, 3'-amidates, MMI, alkyl ether (U. S. Patent No. 5,223,618) and others as described in U. S. Patent Nos. 5,378,825, 5,489,677, 5,541,307, and the like. Where RNase activity is desired, a backbone capable of serving as an RNase substrate is employed for at least a portion of the oligonucleotide.

Universal bases suitable for use in the present invention include, but are not limited to, deoxy 5-nitroindole, deoxy 3-nitropyrrole, deoxy 4-nitrobenzimidazole, deoxy nebularine, deoxyinosine, 2'-OMe inosine, 2'-OMe 5-nitroindole, 2'-OMe 3-nitropyrrole, 2'-F inosine, 2'-F nebularine, 2'-F 5-nitroindole, 2'-F 4-nitrobenzimidazole, 2'-F 3-nitropyrrole, PNA-5-nitroindole, PNA-nebularine, PNA-inosine, PNA-4-nitrobenzimidazole, PNA-3-nitropyrrole, morpholino-5-nitroindole, morpholino-nebularine, morpholino-inosine, morpholino-4-nitrobenzimidazole, morpholino-3-nitropyrrole,

phosphoramidate-5-nitroindole, phosphoramidate-nebularine, phosphoramidate-inosine, phosphoramidate-4-nitrobenzimidazole, phosphoramidate-3-nitropyrrole, 2'-O-methoxyethyl inosine, 2'-O-methoxyethyl nebularine, 2'-O-methoxyethyl 5-nitroindole, 2'-O-methoxyethyl 4-nitro-benzimidazole, 2'-O-methoxyethyl 3-nitropyrrole, deoxy R, MP-5-nitroindole dimer 2'-OMe R, MP-5-nitroindole dimer and the like.

5 Degenerate bases suitable for use in the present invention include, but are not limited to, deoxy P (A&G), deoxy K (U&C), 2'-OMe 2-aminopurine (U&C), 2'-OMe P (G&A), 2'-OMe K (U&C), 2'-F-2-aminopurine (U&C), 2'-F P (G&A), 2'-F K (U&C), PNA-2-aminopurine (U&C), PNA-P (G&A), PNA-K (U&C), morpholino-2-aminopurine (U&C), morpholino-P (G&A), morpholino-K (U&C), phosphoramidate-2-aminopurine (C&U), phosphoramidate-P (G&A), phosphoramidate-K (U&C), 2'-O-methoxyethyl 2-aminopurine (U&C), 2'-O-methoxyethyl P (G&A), 2'-O-methoxyethyl K (U&C), deoxy R, MP-KP dimer, deoxy R, MP-PK dimer, deoxy R, MP-Kk dimer, deoxy R, MP-PP dimer, 2'-OMe R, MP-KP dimer, 2'-OMe R, MP-PK dimer, 2'-OMe R, MP-KK dimer, 2'-OMe R, MP-PP dimer and the like.

10 The present invention provides methods for use of universal and/or degenerate bases in antisense oligonucleotides to provide single antisense molecules that target more than one gene. These universal and/or degenerated bases can be used in either the RNase H portion or non-RNase H portion of antisense molecules. The ability to bind to more than one base on a target provides the flexibility of making one antisense molecule that targets more than one RNA sequence.

Oligonucleotide synthesis is well known in the art, as is synthesis of oligonucleotides containing modified bases and backbone linkages. In one embodiment of the present invention, there is provided an antisense phosphorothioate oligonucleotide having between 6 and about 50 bases in which at least one of its bases are replaced with universal and/or degenerate bases. In a preferred embodiment, no more than about 50% of the bases are universal and/or degenerate bases. Another oligonucleotide for use in the present invention comprises a non-RNase recruiting portion of between 3 and about 15 bases, followed by an RNase-recruiting portion of between 3 and about 15 bases, followed by a second non-RNase H-recruiting portion of 3 to about 15 bases, wherein at least one of the bases contained in the oligonucleotide are degenerate and/or universal bases. In a preferred embodiment, no more than about 50% of the bases are universal and/or degenerate bases. Another antisense oligonucleotide contemplated for use in the present invention comprises a non-P¹ase H recruiting portion followed by a RNase H-recruiting portion in which at least one of its bases are replaced with universal and/or degenerate bases. In a preferred embodiment, no more than about 50% of the bases are universal and/or degenerate bases. An antisense oligonucleotide comprising an RNase H-recruiting portion followed by a non-RNase H-recruiting portion, in which at least one of its bases are replaced with degenerate and/or universal bases, is also within the scope of the present invention. In a preferred embodiment, no more than about 50% of the bases are universal and/or degenerate bases.

Other antisense oligonucleotides contemplated for use in the present invention include: an oligonucleotide comprising an RNase L recruiting oligonucleotide 2'-5' adenosine moiety in which the oligonucleotide comprises at least one degenerate and/or universal base; and an oligonucleotide designed to recruit RNase P in which the oligonucleotide

comprises at least one degenerate and/or universal base. In a preferred embodiment, no more than about 50% of the bases are universal and/or degenerate bases.

Another embodiment of the invention is a ribozyme in which at least one base within the RNA targeting sequence is a degenerate and/or universal base. In a preferred embodiment, no more than about 50% of the bases are universal and/or degenerate bases. The minimum sequence requirements for ribozyme activity are described by Bensele et al. (*J. Am. Chem. Soc.* 115:8483-8484, 1993). Hammerhead ribozyme molecules comprise end domains ("I" and "III") which hybridize to the substrate polynucleotide, a catalytic portion, and a stem loop structure ("II") which can be substituted by a variety of other structures capable of holding the molecule together.

The antisense oligonucleotides of the present invention can be used to target one or more genes, more preferably therapeutic genes, and most preferably anti-apoptosis or chemoresistance genes as described in the examples presented below.

Representative classes of antisense oligonucleotides for use in the present invention are shown below. Although this figure shows 18-mers, this should be considered illustrative rather than limiting.

- 15 5'-NNN NNN BBB BBB NNN NNN-3' (SEQ ID NO: 1)
 5'-NNN NNN BBB BBB NNN NNN-3' (SEQ ID NO: 2)
 5'-NNN NNN BBB BBB NNN NNN-3' (SEQ ID NO: 3)
 5'-NNN NNN BBB BBB NNN NNN-3' (SEQ ID NO: 4)
 5'-NNN BNN BBN BNB NBN NBN-3' (SEQ ID NO: 5)
 20 5'-NNN BNN BBN BNB NBN NBN-3' (SEQ ID NO: 6)
 5'-NNN BNN BBN BNB NBN NBN-3' (SEQ ID NO: 7)
 5'-a⁺a⁺a⁺a⁺----NNN BNN BBN BNB NBN NBN-3' (SEQ ID NO: 8)
 5'-NNN BNN BBN#BNB NBN NBN-3' (SEQ ID NO: 9)
 5'-NNN BNN BBN&BNB NBN NBN-3' (SEQ ID NO: 10)
 25 5'-NNN BNN BBN BNB NBN NBN-3' (SEQ ID NO: 11)

In these sequences, B is a universal base or degenerate base; N is a natural or non-naturally occurring base capable of specific recognition of an RNA target base including, but not limited to, A, C, G, T, U, propynyl C, propynyl U, diamopurine, 5-MeC, 7-deaza A and 7-deaza G. The underline represents the non-RNase H recruiting section, including, but not limited to, 2'-O-alkyl, PNA, methylphosphonate, 3'-amidate, 2'-F, morpholino, 2'-O-alkylaminoalkyl and 2'-alkoxyalkyl. The "----" represents a linker including, but not limited to the one disclosed in U. S. Patent No. 5,583,032. The "#" represents the ribozyme cleaving portion of a ribozyme oligonucleotide; the "&" represents the stem loop structure that recruits RNase P; and a⁺a⁺a⁺a⁺ represents a tetramer of oligomeric 2'-5' adenosine. SEQ ID NO: 11 is also designed to recruit RNase P by inducing formation of a loop structure on the target RNA which is a substrate for RNase P (See U.S. Patent No. 5,877,162).

The antisense oligonucleotides and ribozymes described above are used to cleave one or more target RNA molecules *in vitro* or *in vivo*.

Example 1

Oligonucleotide synthesis

5 All reagents are used dry (<30 ppm water). Oligonucleotide synthesis reagents are purchased from Glen Research. Amidites in solution are dried over Trap-paks (Perkin-Elmer Applied Biosystems, Norwalk, CT). A solid support previously derivatized with a dimethoxy trityl (DMT) group protected propyl linker is placed in a DNA synthesizer column compatible with a Perkin-Elmer Applied Biosystems Expedite synthesizer (1 mmol of starting propyl linker). The DMT group is removed with a deblock reagent (2.5% dichloroacetic acid in dichloromethane). The
10 standard protocols for RNA and DNA synthesis are applied to amidites (0.1 M in dry acetonitrile). The amidites are activated with tetrazole (0.45 M in dry acetonitrile). Coupling times are typically up to 15 minutes depending on the amidite. The phosphonite intermediate is treated with an oxidizing Beaucage sulfurizing reagent. After each oxidation step, a capping step is performed which places an acetyl group on any remaining uncoupled 5'-OH groups by treatment with a mixture of two capping reagents: CAP A (acetic anhydride) and CAP B (n-methylimidazole in THF). The cycle is
15 repeated a sufficient number of times with various amidites to obtain the desired sequence. After the desired sequence is obtained, the support is treated at 55°C in concentrated ammonium hydroxide for 16 hours. The solution is concentrated on a speed vac and the residue is taken up in 100 ml aqueous 0.1 M triethylammonium acetate. This is applied to an HPLC column (C-18, Kromasil, 5 mm, 4.3 mm diameter, 250 mm length) and eluted with an acetonitrile gradient (solvent A, 0.1 M TEAA; solvent B, 0.1 M TEAA and 50% acetonitrile) over 30 minutes at 1 ml/min flow rate.
20 Fractions containing greater than 80% pure product are pooled and concentrated. The resulting residue is taken up in 80% acetic acid in water to remove the trityl group and reapplied to a reverse phase column and purified as described above. Fractions containing greater than 90% purity are pooled and concentrated.

The antisense activity of the oligonucleotides of the invention can be determined by standard assay methods as described, for example, in Examples 2-4. In general, one can prepare a target polynucleotide having a known
25 sequence, contact the target with oligomers of the invention selected to bind the target sequence to form a complex, subject the complex to cleavage with the desired target nuclease and analyze the products to determine if cleavage occurred. The activity can be determined by detecting cleaved target polynucleotides directly (e.g., by hybridization to a labeled probe, amplification by PCR, visualization on a gel, and the like), or by an effect on a host cell phenotype (for example, expression or lack of expression of a selected protein). The RNase H cleavage assay is described below

30

Example 2

RNase H cleavage assay

PCR is used to prepare a dsDNA fragment encoding part of secreted alkaline phosphatase (SEAP) using the following primers:

P3 - 5'-CGAAA-TTAAATCGACTCACTAT-3' (SEQ ID NO: 12),

P3.1 - 3'-GCTTTAATTATGCTGAGTGATATCCCGAAGCTTAGCGCTTAAGCGGGTGGT-

ACGACGACGACGACGACGACGACCCGGAC-5' (SEQ ID NO: 13);

P4 - 3'-TAGGGTCAACTCCTCCTTGG-5' (SEQ ID NO: 14); and

P5 - 3'-TACGAC-GACGACGACGACGACGACCCGGACTCCGATGTCGAGAGGGACCCGTAGTA-

5 GGGTCAACTCCTCCTTGG-5' (SEQ ID NO: 15).

These primers are based on the SEAP RNA fragment (1 to 102) having the sequence: 5'-GGGCTTCGAATCGCGAATTCGCCACCATGCTGCTGCTGCTGCTGGGCTGAGGCTACAGCTCTCCCTGGGCATCATCCCAGTTGAGGAGGAGAACC-3' (SEQ ID NO: 16).

10 PCR amplification is performed under the manufacturer's (Life Technologies) recommendation reaction conditions. Primers P3.1 and P5 are used at 10 nM, while primers P3 and P4 are used at 0.50 μ M. The PCR program is 94°C for 5 minutes, 35 cycles at 52°C for 30 seconds, 72°C for 1 minute, 94°C for 45 seconds and 72°C for 10 minutes.

SEAP dsDNA is then transcribed into ssRNA using a RiboMax™ large scale RNA kit (Promega, Madison, WI). The SEAP DNA concentration is 30 μ g/ml. The transcription reaction is terminated by adding DNase I and incubating
15 at 37°C for 15 minutes. DNA fragments and free nucleotides are removed by precipitation in ethanol/sodium acetate and washing with 70% ethanol. The RNA was suspended and diluted to approximately 2 μ M for use in the RNase H activity assays.

Oligonucleotides of the present invention complementary to a portion of SEAP RNA (20 μ M each), SEAP RNA (10 μ l of 2 μ M solution), and Tris/EDTA buffer (10 mM Tris-HCl, pH 7.4, 1 mM EDTA, "TE", qs to 2 μ l) are
20 added to 500 μ l thin-wall reaction tubes and incubated for 3 to 5 minutes at 40°C to reach thermal equilibrium. RNase H buffer (10X: 200 mM Tris-HCl, pH 7.4-7.5, 1,000 mM KCl, 100 mM MgCl₂·6H₂O, 0.5 mM dithiothreitol, 25% w/v sucrose), RNase H (0.4 to 0.6 U, Promega), and water (qs to 20 μ l), are combined to form a cocktail, and incubated for 3 to 5 minutes at 40 °C. Then, 8 μ l of the cocktail is added to each reaction tube and mixed as quickly as possible to prevent cooling. Reactions are incubated at 40°C for 30 minutes in an MJ Research (Watertown, MA)
25 PCT-100 temperature controller. Reactions are stopped by adding 20 μ l FDE sample buffer (90% v/v formamide, 10% v/v 10X TBE buffer, 0.5% w/v bromphenol blue, 25 mM EDTA) (1XTBE: 89 mM Tris base, 89 mM boric acid, 2 mM EDTA, pH 8.0) to each reaction and heating to 90°C for 3 to 5 minutes.

Each sample (8 to 10 μ l) is subjected to polyacrylamide gel electrophoresis on denaturing 15% gels at 200 volts for about one hour, or until the dye front reaches the bottom of the gel. Nucleic acid bands in gels are visualized
30 by soaking the gels in a 1:10,000 dilution of Cyber Gold™ (Molecular Probes, Junction City, OR) in 1X TBE for 5-10 minutes, soaking in 1X TBE for an additional 5-10 minutes and irradiating on a short wave UV transilluminator. The results are recorded by photographing the CyberGold™ fluorescence using a CyberGREEN™ filter and a Polaroid MP-4 camera with Polaroid Type 667 3000 ASA black and white film.

Duplex DNA ladders (20 bp and 100 bp, GenSura, San Diego) are used as size standards. Standard ladders are not heated before loading onto gels, and are undenatured, running as duplex DNA fragments in both denaturing and non-denaturing gels.

Example 3

5 Intracellular antisense activity against protein kinase C alpha (PKC α)

Protein kinase C alpha (PKC α) is used as a gene target to demonstrate antisense activity of the oligonucleotides comprising degenerate and/or universal bases of the invention. PKC α is a normal human gene that is overexpressed in a majority of human cancer types, and is one of the most highly publicized of all antisense target genes.

10 A human bladder carcinoma cell line (T-24, ATCC HTB-4), a cell line known to overexpress PKC α , is cultured using standard methods: 37°C, 5% CO₂ in 75 cm² flasks in McCoy's 5A medium (Mediatech, Herndon, VA) with 10% fetal bovine serum and penicillin-streptomycin. For antisense experiments, T-24 cells are plated into 12-well plates at 75,000 cells/well and allowed to adhere and recover overnight before transfection. The oligonucleotide 5'-GTTCTCXXXXXGAGTTT-3' (SEQ ID NO: 17) in which the X residues are universal and/or degenerate bases (the same
15 or different), and in which remaining residues are connected by modified backbone linkages other than phosphorothioate linkages, and a control oligonucleotide, are transfected into T-24 cells using a cationic lipid-containing cytofection agent (LipofectACE™) (GibcoBRL, Gaithersburg, MD) which provides efficient nuclear delivery of fluorescently labeled oligonucleotides of the invention in T-24. This is an analog of 5'-GTTCTCGCTGGTGAGTTTCA-3' (SEQ ID NO: 18) which is a known PKC α antisense molecule.

20 Oligonucleotides of the invention and conventional all-phosphorothioate oligonucleotides are diluted into 1.5 ml of reduced serum medium Opti-MEM⁺ I (GibcoBRL) to a concentration of 400 nM each. The oligonucleotide-containing solutions are then mixed with an equal volume of Opti-MEM I containing LipofectACE sufficient to give a final lipid to oligonucleotide ratio of 5 to 1 by weight. The final concentration of oligonucleotide is 200 nM. The oligonucleotide/lipid complexes are incubated at room temperature for 20 minutes before adding to tissue culture cells.

25 Cells are washed once in phosphate buffered saline (PBS) to rinse away serum-containing medium, followed by addition of 1 ml transfection mix to each well of a 12-well plate. All transfections are performed in triplicate. The cells are allowed to take up oligonucleotide/lipid complexes for 22 hours prior to harvesting the total cellular RNA. Mock transfections consist of cells treated with Opti-MEM 1 only.

30 After 22 hours of antisense treatment, total RNA is harvested from the cells. The cells are released from the plates by trypsin/EDTA treatment according to standard methods. The triplicate groups of cells are pooled and total cytoplasmic RNA is isolated using an RNeasy kit (QIAGEN) according to the manufacturer's protocols. The RNA is treated with DNase I and UV quantitated according to standard methods.

35 Reverse transcriptase/polymerase chain reaction (RT-PCR) is performed with the methods and materials from a SuperScript One-Step RT-PCR kit from GibcoBRL. The RT-PCR reactions to detect PKC α are performed in two independent runs, with PKC α -specific primers from Oxford Biomedical Research and 100 ng of input total RNA.

Control multiplex RT-PCRs (MP RT-PCRs) are performed to confirm equal quantities of input RNA into each PKC α RT-PCR. The primers, reagents and protocol are from Maxim Biotech. The control MP RT-PCRs amplify BAX and LICE genes equally in all samples, confirming that equal amounts of intact RNA are added to the PKC α RT-PCRs.

All RT-PCR reactions are performed according to the following program of a PTC-1000 thermocycler (MJ Research): Step 1, 50°C for 35 minutes; Step 2, 94°C for 2 minutes; Step 3, 55°C for 30 seconds; Step 4, 72°C for 1 minute; Step 5, 94°C for 30 seconds; Step 6, go to step 3, 33 more times; Step 7, 72°C for 10 minutes; Step 8, end. all RT-PCR products are separated on a 4% Super Resolution Agarose TBE gel (Apex) and stained with Cyber Gold™ according to the manufacturer's instructions. Gels are photographed on Polaroid Type 667 film.

Example 4

10 Antisense activity against human Bcl2 gene in tissue culture cells

B cell lymphoma-associated gene 2 (Bcl2) is a "normal" human gene that is overexpressed in a majority of human cancer types. The Bcl2 protein regulates cell death and BCL overexpression is known to cause cells to be chemotherapy and radiation resistant. The following Bcl2-targeted antisense molecule is synthesized:

5'-TCTXCCXXCXTXCCXT-3' (SEQ ID NO: 19), in which X is the same or different universal and/or degenerate
15 bases, and in which the first nine residues are a non-RNase H recruiting region (i. e., contain modified backbone linkages other than phosphorothioate linkages). This is an analog of the oligonucleotide
5'-TCTCCAGCGTGCGCCAT-3' (SEQ ID NO: 20).

T-24 cells are plated at 75,000 cells/well and allowed to adhere and recover overnight before oligonucleotide transfections. Test and control oligonucleotides are transfected into T-24 cells using LipofectACE™. Oligonucleotides
20 are diluted into 1.5 ml of reduced serum medium (OptiMEM™, GibcoBRL) to a concentration of 400 nM each. The oligonucleotide-containing solutions are then mixed with an equal volume of Opti-MEM I containing LipofectACE sufficient to give a final lipid to oligonucleotide ratio of 5 to 1 by weight. the final concentration of oligonucleotide is 200 nM. The oligonucleotide/lipid complexes are incubated at room temperature for 20 minutes before adding to tissue culture cells. Cells are washed once in PBS, followed by addition of 1 ml of transfection mixed into each well
25 of a 12-well plate. All transfections are performed in triplicate. Cells are allowed to take up oligonucleotide/lipid complexes for 24 hours prior to harvesting of total cellular RNA. Mock transfections consist of cells treated with Opti-MEM I only. Total cytoplasmic RNA is isolated and quantitated as described in Example 3.

RT-PCR is performed as described in Example 3. The RT-PCR reactions to detect bcl-2 are performed with the primers: 5'-GGTGCCACCTGTGGTCCACCTG-3' (SEQ ID NO: 21) and 5'-CTTCACTTGTGGCCAGATAGG-3' (SEQ
30 ID NO: 22) and 1 μ g of input total RNA. Control RT-PCR reactions against β -actin are also performed using the primers 5'-GAGCTGCGTGTGGCTCCGGAGG-3' (SEQ ID NO: 23) and 5'-CGCAGGATGGCATGGGGGCATACCCC-3' (SEQ ID NO: 24) and 0.1 μ g of input total RNA.

All bcl-2 and β -actin RT-PCR reactions are performed according to the following program on a PTC-100 thermocycler (MJ Research): Step 1, 50°C for 35 minutes; Step 2, 94°C for 2 minutes; Step 3, 60°C for 30 seconds;

Step 4, 72°C for 1 minute; Step 5, 94°C for 30 seconds; Step 6, go to step 3, 35 more times; Step 7, 72°C for 10 minutes; Step 8, end.

All RT-PCR products are separated on a 4% Super Resolution Agarose TBE gel and stained with CyberGold™ according to the manufacturer's instructions. Gels are photographed on Polaroid Type 667 film.

5

Example 5

Antisense targeting of bcl-2A and bcl-xL

Many tumors overexpress multiple chemoresistance genes simultaneously, and are thus unlikely to respond to antisense-based therapies against only one specific chemoresistance gene at a time. Knockout of multiple resistance genes with a single antisense oligonucleotide can enhance chemosensitization in resistant tumors. A known
10 example of such simultaneous expression of chemoresistance genes is bcl-2A and bcl-xL which are distinct, but related, transforming oncogenes are overexpressed in many human cancers. Most importantly, the overexpression of both bcl-2 family members has been shown to confer chemoresistance to cells.

Previously reported attempts to knock out both genes simultaneously were based on conventional oligonucleotides that are perfectly complementary to one gene or the other, but not both, and thus have several
15 mismatches and low activity against one of the target genes. Thus, these attempts have relied on non-specific RNase H-dependent activity of long oligonucleotides. In contrast, the use of two or more oligonucleotides, one targeted against each gene, is far more likely to result in toxic effects and to produce non-specific antisense activity.

The present invention provides a single antisense oligonucleotide for simultaneous knockout of two or more genes. For example, bcl-2 and bcl-xL are simultaneously targeted with a single oligonucleotide containing one or more
20 universal and/or degenerate bases targeted to the small region of high nucleotide homology shown in Figure 1. Six representative antisense oligonucleotides containing one or more universal and/or degenerate bases, and the regions to which they hybridize, are shown in Fig. 1. (Human bcl-2 mRNA (HUMBCL2A) - GenBank #M13994; bcl-xL mRNA (HSBCLXL) - GenBank #Z23115) Asterisks indicate mismatches in the region of nucleotide similarity. Base numbers are as defined in GenBank.

25

Example 6

Targeting of two or more related genes

The protein kinase C (PKC) gene family comprises gene products which regulate cell growth by phosphorylating other proteins in response to extracellular signals. Overexpression of PKC genes has been detected in several human tumor types and PKC genes are believed to be potential cancer therapy targets. Despite the similarity
30 of PKC family members at the protein level, the nucleotide sequences can be significantly different. Antisense oligonucleotides including one or more universal or ambiguous bases allows two or more PKC family members to be targeted at the nucleotide level. Figure 2 shows a sequence alignment of homology regions one and two of human PKCα mRNA (HSPKCA1; GenBank #X52479), human PKCθ mRNA (HUMPKCTH; GenBank #L07860) and human PKCδ mRNA (HUMPKCD13X; GenBank #L07860). Representative oligonucleotides for targeting two or three of these
35 PKC family members are shown in Figure 2.

Example 7Targeting two alleles of the same gene

Comparison of allelic variations in an important human oncogene, bcl-2, reveals several single nucleotide polymorphisms (SNPs) within the general human population. Overexpression of any known allele of bcl-2 has been shown to confer chemoresistance in human tumors and is regarded as a poor prognostic indicator. Two or more alleles of the bcl-2 gene can be targeted with single oligonucleotides including one or more universal or degenerated bases without restriction by the occurrence of SNPs. The two regions of human bcl-2B (HUMBCL2B; GenBank #M13995) and human bcl-2C (HUMBCL2C; GenBank #M14745) are shown in Figure 3, as are representative oligonucleotides which target regions of both alleles.

This allows an antisense oligonucleotide gene walk, the evaluation of a series of antisense oligonucleotides distributed throughout the entire length of overlap between the genetic alleles, to be performed without limitation by the occurrence of SNPs. If SNPs could not be included in the regions targeted by antisense oligonucleotides, the gene walk would be far less effective at identifying effective antisense target sites that yield efficient inhibition of gene expression.

15

Example 8Elimination of problematic antisense base sequence motifs

The oligonucleotides flanked by "###" in Figure 3 illustrate another advantage of incorporation of universal and/or degenerate bases into antisense oligonucleotides, namely the elimination of "CG" dinucleotides and tetra-G sequences which can have deleterious effects as previously discussed. Thus, the use of universal and/or degenerate bases eliminates sequence-dependent, non-antisense effects by substituting universal and/or ambiguous bases into problematic sequence motifs. This is also illustrated below:

20

Anti-bcl-2: 3' GGGCCCGTGTGCGGGTA (SEQ ID NO: 25) (tetra-G)

becomes: 3' GGGCCPGTGTGP_KGGTA (SEQ ID NO: 26)

25

Anti-bcl-2: 3'-CGTCTGGGGCCGACGGGGG (SEQ ID NO: 27) (double tetra-G)

becomes: 3'-CGTCTG_KGGCCGACGG_KGG (SEQ ID NO: 28)

Anti-bcl-2: 3'-GGCCGGGGGGGGCCCCG (SEQ ID NO: 29) (highly CG)

becomes: 3'-GGCPGPGGGPG_KCCCC_PG (SEQ ID NO: 30)

30

While particular embodiments of the invention have been described in detail, it will be apparent to those skilled in the art that these embodiments are exemplary rather than limiting, and the true scope of the invention is that defined in the following claims.

WHAT IS CLAIMED IS:

1. An antisense oligonucleotide having at least one non-naturally occurring backbone linkage and having between 6 and about 50 bases, wherein at least one of said bases are universal and/or degenerate bases.
2. The antisense oligonucleotide of Claim 1, wherein no more than about 50% of said bases are universal and/or degenerate bases.
3. An antisense oligonucleotide comprising a first non-RNase H recruiting region having between 3 and about 15 bases, an RNase H recruiting region having between 3 and about 15 bases, and a second non-RNase H recruiting region, wherein at least one of said bases are universal and/or degenerate bases.
4. The antisense oligonucleotide of Claim 3, wherein no more than about 50% of said bases are universal and/or degenerate bases.
5. An antisense oligonucleotide comprising a non-RNase H recruiting section and an RNase H recruiting section, wherein at least one of said bases are universal and/or degenerate bases.
6. The antisense oligonucleotide of Claim 5, wherein no more than about 50% of said bases are universal and/or degenerate bases.
7. An oligonucleotide comprising an RNase L-recruiting region comprising a 2'-5' adenosine oligomer, wherein the RNA targeting region of said oligonucleotide comprises at least one universal and/or degenerate bases.
8. The oligonucleotide of Claim 7, wherein said RNA targeting region comprises no more than about 50% universal and/or degenerate bases.
9. An oligonucleotide designed to recruit RNase P, wherein the RNA targeting region of said oligonucleotide comprises at least one universal and/or degenerate bases.
10. The oligonucleotide of Claim 9, wherein said RNA targeting region comprises no more than about 50% universal and/or degenerate bases.
11. A ribozyme having an RNA targeting region which comprises at least one universal and/or degenerate bases.
12. The ribozyme of Claim 11, wherein said RNA targeting region comprises no more than about 50% universal and/or degenerate bases.
13. A method for cleaving a target RNA molecule, comprising the step of contacting said RNA molecule with an oligonucleotide according to any one of Claims 1-10 in the presence of an RNase capable of cleaving said target.
14. The method of Claim 13, wherein said RNase is selected from the group consisting of RNase H, RNase L and RNase P.
15. A method for cleaving a target RNA molecule, comprising the step of contacting said RNA molecule with a ribozyme according to Claims 11 or 12.

16. A method for cleaving one or more target RNA molecules, comprising the step of contacting said RNA molecule with an oligonucleotide having between 6 and about 50 bases, wherein said oligonucleotide comprises at least one universal and/or degenerate base.,

17. A method for reducing the deleterious effects of an antisense oligonucleotide comprising one or more sequence motifs, comprising replacing one or more bases within said one or more sequence motifs with one or more universal and/or degenerate bases.

18. The method of Claim 17, wherein said sequence motif is a CG dinucleotide.

19. The method of Claim 17, wherein said sequence motif is a poly-G sequence.

FIG. 1

bcl-2A: 2020 tggatccaggataacggaggctgggatgcctttgtggaact 2060
|||||*|||||*|||||*|*|||||
bcl-xL: 675 tggatccaggagaacggcggtgggatactttgtggaact 715
Examples: 3' ACCTAGGTCCTBTTGCCBCC
3' GGTCTCTBTTGCCBCCGACCC
3' CCTBTTGCCBCCGACCCTAPGKAAACACC
3' GCCBCCGACCCTAPGKAAAC
3' BCCGACCCTAPGK
3' ACCCTAPGKAAACACCTTGA

FIG. 2

Homology region One (PKCa, PKCt, PKCd):

alp : 1276 atggaatatgtcaacgggtggggacctcatgtaccacat 1313
 ||||| | ||||| ||||| ||||| ||||| |||||
 theta: 1393 atggagtacctaacggaggggacttaattgtaccacat 1430
 ||||| | ||||| ||||| ||||| ||||| |||||
 delta: 1337 atggagttcctcaacgggggggacctgatgtac 1369
 Examples: 3' CCTPABKBAGTTGCCBCCC
 3' GCCBCCCCTGKABTACATG

Homology region Two. (PKCa, PKCt, PKCd):

alpha: 1398 aggaatcatttatagggatctgaagttagataaacgtcatgttggattcagaaggacatat 1457
 || || || || ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 theta: 1515 aggaatagtctacagggacctgaagctagataaacatcctgttagacaaagatggacatat 1574
 || || || || ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 delta: 1459 gggcatcatttacaggggacctcaaaactggacaatgtgctgttggacogggatggccacat 1518
 Examples: 3TCCCTKGABTTPKAKCTKT 3' ACAAPCTKBBPCTBCCBGT

alpha: 1458 caaaattgctgactttgggatgtgcaaggaaacacatgatggatggagtcacgaccaggac 1517
 ||| || || || ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 theta: 1575 caagatcgcggaattttggaatgtgcaaggagaacatgttaggagatgccaagacgaatac 1634
 ||| || || || ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 delta: 1519 caagattgcogactttgggatgtgcaagagaacatattcggggagagccggggccagcac 1578
 Examples: 3' CGBCTKAAACCTACACG
 3' PTACACGTTPTPTGTGA
 3' CACGTTPTPTGTAPB

alpha: 1518 cttctgtgggactccagattatatcgccccagagat 1553
 ||||| || || || || ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 theta: 1635 cttctgtgggacacctgactacatcgccccagagat 1670
 ||||| || || || || ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
 delta: 1550 cttctgcggcacccctgactatatcgccccagagat 1614
 Examples: 3' GAAGACKCCBTGEGGACT
 3' KATKTAGCGGRGCTCT

FIG. 3

bcl-2B: 116 gttggccccggttgccttttctctgggaaggatggcgacgctgggagaaacgggggtacga 175
 |||||
 bcl-2C: 1 gttggccccggttacttttctctgggaaatatggcgacgctgggagaaacgggggtacga 60
 Examples: 3' GGGCAAPGAAAAGGAGAC
 3' CCTTPBTACCGCGTCCGACCC
 3' CCCTCTTGPCCCATGCT
 3' CGA-

bcl-2B: 176 caaccgggagatagatgaagtacatccattataagctgtcgcagagggggtacgagtg 235
 |||||
 bcl-2C: 61 taaccgggagatagtgatgaagtacatccattataagctgtcgcagagggggtacgagtg 120
 Examples: -KTTGGCCCTCTATCAC

bcl-2B: 236 ggatgcgggagatgtggcgccgcgccccggggcgcccccgccacggggcatcttctc 295
 |||||
 bcl-2C: 121 ggatgcgggagatgtggcgccgcgccccggggcgcccccgccacggggcatcttctc 180
 Examples: 3' GGGCGPGGCCCGTAGAAG
 3' GAG-

bcl-2B: 296 ctcccagcccgggcacacgccccatccagccgcatcccgcgacccgggtcgccaggacctc 355
 |||||
 bcl-2C: 181 ctgcgagcccgggcacacgccccatacagccgcatcccgggacccgggtcgccaggacctc 240
 Examples: -GAGBGTCCGGCCCGT
 ###3' GGGCCPGTCTGTPGKGGTA###
 3' GGGTABGTCCGGCGTAGG
 3' CGTAGGGCBCTGGGCC

bcl-2B: 356 gccgctgcagaccccggtgccccggcgccgcgccccgggctgcgctcagcccggtgcc 415
 |||||
 bcl-2C: 241 gccgctgcagaccccggtgccccggcgccgcgccccgggctgcgctcagcccggtgcc 300
 Example: . ###3' CGTCTGKGGCCGACGGKGG###
 ###3' GGCPTGGPGGPGGPGCCCPG###

bcl-2B: 416 acctgtggtccacctggccctccgccaagccggcgacgacttctcccgccgctaccgccc 475
 |||||
 bcl-2C: 301 acctgtggtccacctgacccctccgccaagccggcgacgacttctcccgccgctaccgccc 360
 Examples: 3' AGGTGGACPGGGAGGC
 3' GCGCGTTCGCGCCGC
 3' GCGCB-

bcl-2B: 476 cgacttcgcccagatgtccagccagctgcacctgacgcccctcaccgcccggggagcgtt 535
 |||||
 bcl-2C: 361 cgacttcgcccagatgtccagccagctgcacctgacgcccctcaccgcccggggagcgtt 420
 Examples: -CGTGAGCGGCTC
 3' CTGGTCBGTGACGTGG

SEQUENCE LISTING

<110> OASIS BIOSCIENCES, INC.
Brown, Bob D.
Riley, Timothy A.

<120> ANTISENSE OLIGONUCLEOTIDES COMPRISING
UNIVERSAL AND/OR DEGENERATE BASES

<130> OASBIO.001VPC

<150> US 60/128,377

<151> 1999-04-08

<160> 30

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide primers

<221> misc_feature

<222> 1-6, 13-18

<223> n = Natural or non-naturally occurring base capable
of specific recognition of an RNA target base
including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature

<222> 7-12

<223> b = Universal or degenerate base

<400> 1

nnnnnnbbbb bnnnnnnn

18

<210> 2

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide primers

<221> misc_feature

<222> 1-6 13-18

<223> n = Natural or non-naturally occurring base capable
of specific recognition of an RNA target base

including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature
<222> 7-12
<223> b = Universal or degenerate base

<400> 2
nnnnnnbbbbb bnnnnnnn

18

<210> 3
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 1-6, 13-18
<223> Natural or non-naturally occurring base capable of
specific recognition of an RNA target base
including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature
<222> 7-12
<223> b = universal or degenerate base

<400> 3
nnnnnnbbbbb bnnnnnnn

18

<210> 4
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 1-6, 13-18
<223> n = Natural or non-naturally occurring base capable
of specific recognition of an RNA target base
including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature
<222> 7-12
<223> b = Universal or degenerate base

<400> 4

nnnnnnbbbb bnnnnnnn

18

<210> 5
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 1-3, 5-6, 9, 11, 13, 15-16, 18
<223> n = Natural or non-naturally occurring base capable
of specific recognition of an RNA target base
including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature
<222> 4, 7-8, 10, 12, 14, 17
<223> b = Universal or degenerate base

<400> 5
nnnbnnbbnb nbnbnbn

18

<210> 6
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 1-3, 5-6, 9, 11, 13, 15-16, 18
<223> n = Natural or non-naturally occurring base capable
of specific recognition of an RNA target base
including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature
<222> 4, 7-8, 10, 12, 14, 17
<223> b = Universal or degenerate base

<400> 6
nnnbnnbbnb nbnbnbn

18

<210> 7
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 1-3, 5-6, 9, 11, 13, 15-16, 18
<223> n = Natural or non-naturally occurring base capable
of specific recognition of an RNA target base
including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature
<222> 4, 7-8, 10, 12, 14, 17
<223> b = Universal or degenerate base

<400> 7
nnnbnbbbnb nbnbnbn

18

<210> 8
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 1-3, 5-6, 9, 11, 13, 15-16, 18
<223> n = Natural or non-naturally occurring base capable
of specific recognition of an RNA target base
including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature
<222> 4, 7-8, 10, 12, 14, 17
<223> b = Universal or degenerate base

<400> 8
nnnbnbbbnb nbnbnbn

18

<210> 9
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 1-3, 5-6, 9, 11, 13, 15-16, 18
<223> n = Natural or non-naturally occurring base capable
of specific recognition of an RNA target base
including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature
<222> 4, 7-8, 10, 12, 14, 17
<223> b = Universal or degenerate base

<400> 9
nnnbnbbbnb nbnbnbn

18

<210> 10
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 1-3, 5-6, 9, 11, 13, 15-16, 18
<223> n = Natural or non-naturally occurring base capable
of specific recognition of an RNA target base
including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature
<222> 4, 7-8, 10, 12, 14, 17
<223> b = Universal or degenerate base

<400> 10
nnnbnbbbnb nbnbnbn

18

<210> 11
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 1-3, 5-6, 9, 11, 13, 15-16, 18
<223> n = Natural or non-naturally occurring base capable
of specific recognition of an RNA target base
including, but not limited to, a, c, g, t, u,
propynyl C, propynyl U, diaminopurine, 5-MeC,
7-deaza A and 7-deaza G

<221> misc_feature
<222> 4, 7-8, 10, 12, 14, 17
<223> b = Universal or degenerate base

<400> 11
nnnbnbbbnb nbnbnbn

18

<210> 12
<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide primers

<400> 12

cgaaattaaa tcgactcact at

22

<210> 13

<211> 80

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide primers

<400> 13

caggcccagc agcagcagca gcagcagcat ggtgggcgaa ttgcgattc gaagccctat 60
agtgcgtcgt attaatctcg 80

<210> 14

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide primers

<400> 14

ggttctcctc ctcaactggg at

22

<210> 15

<211> 76

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide primers

<400> 15

ggttctcctc ctcaactggg atgatgccca gggagagctg tagcctcagg ccagcagca 60
gcagcagcag cagcat 76

<210> 16

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide primers

<400> 16

gggcttcgaa tcgcgaattc gccaccatg ctgctgctgc tgctgctggg cctgaggcta 60
cagctctccc tgggcatcat ccagttgag gaggagaacc 100

<210> 17
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 7-12
<223> b = Universal or degenerate base

<400> 17
gttctcbbbb bbgagttt 18

<210> 18
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<400> 18
gttctcgtg gtgagtttca 20

<210> 19
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 4, 7-8, 10, 12, 14, 17
<223> b = Universal or degenerate base

<400> 19
tctbccbbcb tbcbbcbt 18

<210> 20
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<400> 20
tctcccagcg tgcgcat 18

<210> 21
<211> 22

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<400> 21
ggtgccacct gtggtccacc tg 22

<210> 22
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<400> 22
cttcacttgt ggcccagata gg 22

<210> 23
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<400> 23
gagctgctgt tggctcccga gg 22

<210> 24
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<400> 24
cgcaggatgg catggggggc ataccc 26

<210> 25
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<400> 25
gggcccgtgt gcggggta 18

<210> 26
<211> 18

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 6, 12
<223> n = 6H, 8H-3, 4-dihydropyrimido [4,5-c] [1,2]
oxazin-7-one

<221> misc_feature
<222> 14
<223> n = 2-amino-6-methoxy aminopurine

<400> 26
gggccngtgt gngnggta 18

<210> 27
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<400> 27
cgtctggggc cgacggggg 19

<210> 28
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<221> misc_feature
<222> 7, 17
<223> k = 2-amino-6-methoxy aminopurine

<400> 28
cgtctgkggc cgacggkgg 19

<210> 29
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic oligonucleotide primers

<400> 29
ggccgcggcg gcgccccg 18

<210> 30
 <211> 18
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Synthetic oligonucleotide primers

 <221> misc_feature
 <222> 4, 6, 9, 12, 17
 <223> n = 6H, 8H-3, 4-dihydropyrimido [4,5-c] [1,2]
 oxazin-7-one

 <400> 30
 ggcnngggng gngccnng

18

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/09293

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : C12Q 1/68; C12P 19/34; C12N 15/11, 15/85; C07H 21/04 US CL : 435/6, 91.3, 325, 366, 371, 375; 536/23.1, 24.3, 24.33, 24.5 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/6, 91.3, 325, 366, 371, 375; 536/23.1, 24.3, 24.33, 24.5 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Extra Sheet.		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 99-18238 A1 (OASIS BIOSCIENCES, INC.) 15 April 1999, see pages 2-3.	1-19
X	REYNOLDS et al. Antisense oligonucleotides containing an internal, non-nucleotide-based linker promote site-specific cleavage of RNA. Nucleic Acid Research. 15 February 1996, Vol. 24, No. 4, pages 760-765, especially page 760.	1-8
X	KRUPP G. Antisense oligoribonucleotides and RNase P. A great potential. Biochimie. May 1993, Vol. 75, No. 1/2, pages 135-139, especially page 138, Figure 2.	9-10
—		—
Y		11-16
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *B* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *Z* document member of the same patent family		
Date of the actual completion of the international search 09 JUNE 2000		Date of mailing of the international search report 08 AUG 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer JANET LEE EPPS Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/09293

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

CAPLUS, BIOSIS, MEDLINE, USPAT, EPO, JPO, DERWENT

search terms: antisense or aptamer or triplex or oligonucleotide or ribozyme, degenerate or universal, RNase P or RNase H or RNase L.